

Final Report for ONR Grant N00014-90-J-1206

§1. Introduction.


This final report covers the time period from December 1, 1989 to November 30, 1995 of ONR Grant N00014-90-J-1206. It is organized as follows. In this section I will give statistical information covering this period. The next five sections progress in the major areas of my research is summarized and the main accomplishments are highlighted. The last section provides references to my papers and other papers discussed in the summary. It is assumed that the reader is familiar with my original proposals [64] and [65].

The papers [27] - [43], [45] - [49], [51], [53], [56], and [57] were published during this period. The papers [44], [50], [52], [55], and [58] were accepted for publication. The papers [54] and [60] were submitted for publication. The papers [29], [39], [41] - [44], [53], and [60] are about on-line coloring. The papers [37], [45], and [54] concern the closely related subject of game chromatic number. The papers [33], [35] - [36], [38], [46], [50], and [57] deal with Gyárfás' graph coloring conjecture on which many of the on-line coloring results rest. The papers [28], [30], [34], [47], [49], [52], and [58] concern the dimension of ordered sets, while [27], [31], [32], and [40] deal with other concepts related to ordered sets. Finally [48], [51], [55], and [56] concern a conjecture of Pósa about square cycles in dense graphs.

The following students, who all received some support from the grant, have all earned doctorate degrees from Arizona State University: Stephen Penrice, Jun Qin, Katalin Kolossa, Juan Quintana, and Yingxing Zhu.

§2. Gyárfás' Conjecture.

Much of my work on on-line coloring is founded on the study of off-line coloring, and in particular, progress on the following conjecture of Gyárfás. For two graphs G and H , let $\text{Forb}(G, H)$ denote the class of graphs that induce neither G nor H . Let $\chi(G)$ denote



19971014 037

DTIC QUALITY INSPECTED 3

ASU
ARIZONA STATE UNIVERSITY

October 2, 1997

Defense Technical Information Center
Building 5, Cameron Station
Alexandria, Virginia 22314

Subject: Grant N00014-90-J-1206
ASU TKA 1761

Enclosed are four copies of the Final Technical Report for the referenced grant, which was under the direction of Dr. Henry A. Kierstead, Department of Mathematics.

If you have questions, please contact Joe Wessels, Sponsored Projects Officer, at (602) 965-1427.

Janice Cox

Janice
Janice D. Bennett
Director

jm

Enclosures

xc: Dr. Henry A. Kierstead

denote the chromatic number of G and $\omega(G)$ denote the clique number of G . Gyárfás made the following conjecture.

Conjecture 2.1. (Gyárfás [68]) For any tree T and clique K , there exists a bound b such that for every graph $G \in \text{Forb}(T, K)$, $\chi(G) \leq b$.

Gyárfás [69] gave easy arguments to show that the conjecture is true if T is a path or a star. Gyárfás, Szemerédi, and Tuza [70] showed that the conjecture is true if T is a radius two tree and K is a triangle. My student Steve Penrice and I made the following major breakthrough.

Theorem 2.2. (Kierstead and Penrice[35]) For every radius two tree T and every clique K , there exists a bound b such that for every graph $G \in \text{Forb}(T, K)$, $\chi(G) \leq b$.

This result depended on the notion of template first used in [33]. Over a series of papers [36], [38], [46], [50], and [57] with various coauthors from the list Penrice, Rödl, Trotter, and Zhu, I sharpened this technique to deal with trees of larger radius. With my student Yingxian Zhu I proved the following generalization of Theorem 2.3, which is exciting because the same type of preliminary result [33] in the radius two case led to the proof of Theorem 2.2.

Theorem 2.3. (Kierstead and Zhu [57]) Let T be the radius three tree such that every vertex adjacent to the root has degree two. Then Conjecture 2.1 holds for T .

An important related question has been raised by Sauer. A class of graphs Γ is said to be *vertex Ramsey* if for every $G \in \Gamma$ there exists $H \in \Gamma$ such that if the vertices of H are partitioned into two sets inducing subgraphs H_1 and H_2 , then either H_1 or H_2 contains an induced copy of G .

Conjecture 2.4. (Sauer [71]) The class $\text{Forb}(T, K)$ is not Ramsey for all trees T and cliques K .

It is easy to show that Conjecture 2.1 implies Conjecture 2.5. What makes Conjecture 2.5 especially interesting from my perspective, is that the same template techniques that have been useful for dealing with special cases of Conjecture 2.1 are also useful for more general cases of Conjecture 2.5. Moreover, these more general cases of Conjecture 2.5 have led to the development of more sophisticated versions of the template technique. I believe that it is likely that we will be able to use these more sophisticated versions to make further progress on Conjecture 2.1. Here are some examples. A *spider with toes* is a tree such that the nonleaf vertices form the subdivision of a star. Let $T(k, r)$ be the tree such that the distance between the root and any leaf is r and every nonleaf has k sons. The following two theorems give the only examples of trees for which Conjecture 2.5 is known to hold, but Conjecture 2.1 is not known to hold. In some sense, a spider with toes is a thin tree, while $T(k, r)$ is a full fat tree when k is large. (Of course, $T(1, r)$ is a spider with toes.)

Theorem 2.6. (Kierstead [50]) For every spider with toes T and clique K , $\text{Forb}(T, K)$ is not vertex Ramsey.

Theorem 2.7 (Kierstead and Zhu [57]) For every positive integer r and sufficiently large integer k and every clique K , $\text{Forb}(T(k, r), K)$ is not vertex Ramsey.

I believe that we are now closing in on the techniques necessary to solve the following problem in particular and possibly all of Conjecture 2.1.

Problem 2.8. Prove Conjecture 2.1 for the case of radius three trees.

§3. On-line Coloring

On-line coloring has been the major focus of this project and I have proved four especially important results. The first of these concerns the application of an on-line coloring algorithm to the off-line problem Dynamic Storage Allocation (DSA). DSA is an NP-Complete off-line problem. In the late sixties an off-line approximation algorithm, based on First-Fit coloring of interval graphs, was suggested for DSA. The performance ratio of this algorithm was twice the competitive ratio of First-Fit. However the best upper bound on the competitive ratio of First-Fit was $O(\omega)$. During the seventies this was improved to $O(\log \omega)$, but the question of whether it was constant remained open until 1986 when I [25] provided a upper bound of 40 for First-Fit coloring of interval graphs, and thus 80 for the performance ratio for DSA. I have now significantly improved this result. I observed that with some care First-Fit could be replaced by a better on-line algorithm for coloring interval graphs. This allowed me to prove the following theorem.

Theorem 3.1. (Kierstead [29]) There exists a polynomial time approximation algorithm for DSA that has a performance ratio of 6.

The intuition behind this algorithm is the following. DSA resembles both Bin Packing and Interval Graph Coloring. There exist orderings for both problems, under which First-Fit performs well, but they are not the same. Thus we order the data so that First-Fit performs well for Bin Packing and use an on-line algorithm for interval graph coloring that is guaranteed to perform reasonably well under any input sequence.

My other on-line coloring results concern on-line coloring per se. The next result is based on Theorem 2.2, but is much harder. It identifies classes of graphs with the property that their on-line chromatic number can be bounded in terms of their clique number.

Theorem 3.2. (Kierstead, Penrice, and Trotter [42]) For every radius two tree T and every clique K , there exists an on-line algorithm A and a bound b such that for every graph $G \in \text{Forb}(T, K)$, $\chi_A(G) \leq b$.

In particular, this answers an old question of Schmerl, Problem 7 of my 1989 ONR proposal [64], who asked whether the on-line chromatic number of co-comparability graphs could be bounded in terms of their clique size. The answer is yes since co-comparability graphs do not induce the radius two tree obtained from a star with three leaves by subdividing each edge.

For any integer $k \geq 2$ the on-line chromatic number of the class of graphs with chromatic number k is unbounded as the number of vertices n increases. Thus in general we must express bounds in terms of n . Lovász, Saks, and Trotter went to considerable effort to prove the upper bound of $O\left(\frac{\log^{(2k-3)} n}{\log^{(2k-4)} n}\right)$, which is barely sublinear. Very recently, I have done much better. I proved the following Theorem, which answers Problem 1 of my 1992 ONR proposal [65].

Theorem 3.3. (Kierstead [60]) There exists an on-line algorithm A that will properly color any k -colorable on-line graph on n vertices with $n^{1-1/k!}$ colors.

My student Katie Kolossa and I had previously obtained much better upper bounds for perfect graphs. Moreover our lower bounds for perfect graphs match the best lower bounds for graphs in general. This answers Problem 2 of my 1992 ONR proposal [65].

Theorem 3.4. (Kierstead and Kolossa [53]) There exists an on-line algorithm A that will properly color any k -colorable on-line perfect graph on n vertices with $n^{10k/\log \log n}$ colors. Moreover for every fixed positive integer k and every on-line algorithm A , there exists an on-line k -colorable perfect graph G such that A uses at least $\frac{1}{2} \left(\frac{\log_3 n}{2k} \right)^{k-1}$ colors on G .

§4. Game Chromatic Number

In [37] Faigle, Kern, Trotter and I developed the notion of game chromatic number of a graph. A game is played between two players A and B on a graph G. The players play by taking turns coloring the vertices of a graph with colors from a fixed set of n colors so that no two adjacent vertices receive the same color. If at some point in this process one of the players cannot play legally, then B wins; otherwise, when the graph is completely colored A wins. The game chromatic number $\chi_g(G)$ of G is the least n such that A has a winning strategy when the game is played on the graph G with n colors. Our main result was that the game chromatic number of a tree is at most four. We also introduced the problem of whether there exists a constant upper bound on the game chromatic number of planar graphs.

This question was answered by Trotter and me. We introduced the notion of a k -admissible ordering of a graph and proved the following two theorems, which together show that every planar graph has game chromatic number at most 33. We denote the least k such that a graph G has a k -admissible ordering by $a(G)$.

Theorem 4.1. (Kierstead and Trotter [45]) For every graph G, $\chi_g(G) \leq \chi(G)a(G) + 1$

Theorem 4.2. (Kierstead and Trotter [45]) For every planar graph G, $a(G) \leq 8$.

Moreover there exist planar graphs with $a(G) = 8$.

Tuza and I generalized the bound on game chromatic number of trees by phrasing it in terms of the treewidth $\tau(G)$ of an arbitrary graph G. This answers Problem 10 of my 1992 ONR proposal [65].

Theorem 4.3. (Kierstead and Tuza [54]) For every graph G, $\tau(G) \leq 6\tau(G) - 2$.

§5. Dimension of Ordered Sets

An important problem in the theory of ordered sets is to bound the dimension of various ordered sets. I have been able to answer several questions that had been open for many years. One such problem concerns the relation between the dimension of an ordered set and the maximum degree Δ of its comparability graph. In a beautiful paper Füredi and Kahn proved the following inequality, where $g(\Delta)$ is the maximum dimension of an ordered set whose comparability graph has maximum degree Δ .

Theorem 5.1. (Füredi and Kahn [67]) For all Δ , $g(\Delta) = O(\Delta \log^2 \Delta)$.

The only problem with this result was that nobody knew of any ordered sets with dimension greater than $\Delta+1$. Erdős, Trotter, and I used random bipartite ordered sets to provide such an example and answer Problem 12 of my 1989 ONR proposal. Let $\Omega(n,p)$ be the probability space consisting of bipartite ordered sets with n maximal elements and n minimal elements where the event that a given maximal element is greater than a given minimal element is independently distributed with probability p .

Theorem 5.2. (Erdős, Kierstead, and Trotter [34]) For every $\varepsilon > 0$ there exists $\delta > 0$ such that if

$$n^{-1+\varepsilon} \leq p \leq \frac{1}{\log n},$$

then for almost all $P \in \Omega(n,p)$, $\dim(P) > \delta \Delta(P) \log |P| > \delta \Delta(P) \log \Delta(P)$.

We also proved:

Theorem 5.3. (Erdős, Kierstead, and Trotter [34]) There exist constants c_1 and c_2 such that for almost all labeled ordered sets P , $\frac{n}{4} \left(1 - \frac{c_1}{\log n}\right) < \dim(P) < \frac{n}{4} \left(1 + \frac{c_2}{\log n}\right)$.

Füredi and Kahn obtained the bound of Theorem 5.1 in terms of the dimension of a certain subset of the boolean lattice. Let B_n denote the Boolean lattice obtained by ordering the

power set of $\{1, 2, \dots, n\}$ by inclusion. Let $B_n(j, k)$ be the restriction of B_n to subsets of size j and k and $\dim(j, k; n) = \dim(B_n(j, k))$. Füredi and Kahn actually proved the following two lemmas.

Lemma 5.4. (Füredi and Kahn [67]) For all ordered sets P with $\Delta(P) = \Delta$,

$$\dim(P) = O\left(\frac{\Delta}{\log \Delta} \dim(1, \log \Delta; \Delta \log \Delta)\right).$$

Lemma 5.5. (Füredi and Kahn [67]) For all positive integers $k < n$, $\dim(1, k; n) \leq (k+1)^2 \log n$. In particular, $\dim(1, \log n; n) = O(\log^3 n)$.

The $\log \Delta$ gap between the example of Theorem 5.2 and Theorem 5.1 made the question of determining $\dim(1, \log n; n)$ important. I proved:

Theorem 5.6. (Kierstead [52]) $\frac{\log^3 n}{64 \log \log n} < \dim(1, \log n; n)$.

In [49] Brightwell, Kostochka, Trotter, and I studied the function $\dim(i, j; n)$ for $1 < i < j$.

§6. Square Cycles in Dense Graphs

An important problem in graph theory is to show that all dense graphs contain certain spanning subgraphs. For example Dirac's famous theorem says that every graph on n vertices with minimum degree at least $n/2$ has a hamiltonian cycle. In 1963 Pósa made the following conjecture.

Conjecture 6.1. (Pósa [66]) Every graph on n vertices with minimum degree at least $\frac{2}{3}n$ contains the square of a hamiltonian cycle, i.e., a hamiltonian cycle together with every 2-chord.

In a series of papers with Genghua Fan, I have come very close to proving this conjecture. First we showed that it was asymptotically correct in [48]. Then we proved the following theorem.

Theorem 6.2. (Fan and Kierstead [51]) Every graph on n vertices with minimum degree at least $(2n-1)/3$ contains the square of a hamiltonian path.

This already implies the important result of Aigner and Brandt that every graph on n vertices with minimal degree at least $(2n-1)/3$ contains every graph on n vertices with maximum degree 2. We also proved:

Theorem 6.3. (Fan and Kierstead [56]) Let G be a graph on n vertices with minimum degree $\frac{2}{3}n$. If G does not contain a square hamiltonian cycle then:

- (1) G does not contain a square cycle of length greater than $\frac{2}{3}n$; and
- (2) The vertices of G can be partitioned into two square cycles.

With my student Juan Quintana [55], I showed that the conjecture is true if G has a maximal 4-clique.

§7. References.

Publications of Henry A. Kierstead

1. "Countable models of ω_1 -categorical theories in admissible languages", **Annals of Math. Logic** **19** (1980), 127-175.
2. "An effective version of Dilworth's theorem", **Trans. Amer. Math. Soc.** **268** (1981), 63-77.
3. "Recursive colorings of highly recursive graphs", **Canadian Journal of Mathematics** **33** (1981), 1279-1290.
4. "An extremal problem in recursive combinatorics", **Congressus Numerantium** **33** (1981), 143-153 (with W. T. Trotter, Jr.).
5. "Indiscernibles and decidable models", **Journal of Symbolic Logic** **48** (1983), 21-32 (with J. B. Remmel).

6. "An effective version of Hall's theorem", **Proc. Amer. Math. Soc.** **88** (1983) 124-128.
7. "Some applications of Vizing's theorem to vertex colorings of graphs" **Discrete Math** **45** (1983), 277-285 (with J. Schmerl).
8. "On the chromatic index of multigraphs without large triangles", **J. of Combinatorial Theory Series B** **36** (1984), 156-160.
9. "A theory of recursive dimension for ordered sets", **Order** **1** (1984), 67-82 (with G. F. McNulty and W. T. Trotter, Jr.).
10. "On coloring graphs with locally small chromatic number", **Combinatorica** **4** (1984) 183-185 (with E. Szemerédi and W. T. Trotter, Jr.).
11. "A new method of proving theorems on chromatic index", **Discrete Math** **52** (1984) 159-164 (with V. Faber and A. Ehrenfeucht).
12. "Degrees of indiscernibles in decidable models", **Trans. Amer. Math. Soc.** **289** (1985), 41-57 (with J. B. Remmel).
13. "Inequalities for greedy dimension of ordered sets", **Order** **2** (1985), 145-164 (with W. T. Trotter, Jr.).
14. "The chromatic number of graphs which induce neither $K_{1,3}$ nor $K_5 - e$ ", **Disc. Math.** **58** (1986), 253-262 (with J. Schmerl).
15. "Recursive ordered sets", **Contemporary Mathematics** **57** (1986), 75-102.
16. "NP-Completeness results concerning greedy and supergreedy linear extensions", **Order** **3** (1986), 123-134.
17. "A Ramsey theoretic problem for ordered sets", **Disc. Math.** **63** (1987) 217-223 (with W. T. Trotter, Jr.).
18. "On π_1^0 -automorphisms of recursive linear orders", **J. Symbolic Logic** **48** (1987), 681-688.
19. "Super greedy linear extensions of ordered sets", **Combinatorial Mathematics: Proceedings of the Third International Conference on Combinatorial Mathematics**, (eds. G. Bloom, R. Graham, and J. Malkevitch), *Annals of the New York Academy of Sciences* **555** (1989), 262-271 (with W. T. Trotter, Jr.).
20. "Hypergraphs with finitely many subtypes", **Trans. Amer. Math. Soc.** **312** (1989), 699-718 (with P. Nyikos).
21. "A minimax theorem for chain complete ordered sets", **Order** **5** (1988), 75-83.
22. "Representing an ordered set as the intersection of super greedy linear extensions", **Order** **4** (1987), 293-311 (with W. T. Trotter, Jr. and B. Zhou).

23. "Explicit matching in middle levels of the Boolean Lattice", **Order** **5** (1988), 163-171 (with W. T. Trotter, Jr.).
24. "Applications of edge coloring of multigraphs to vertex coloring of graphs", **Discrete Math.** **74** (1989), 117-124.
25. "The linearity of First-Fit for coloring interval graphs", **SIAM J. on Discrete Math.** **1** (1988), 526-530.
26. "A proof of Borsuk's theorem", in **Functional Analysis Proceedings, The University of Texas at Austin 1986-1987**, (eds. E. Odell and H. Rosenthal) Springer-Verlag, Berlin (1988), 195-202 (with R. Devore and G. Lorentz).
27. "Racing pawn games", **Congressus Num.** **67** (1988), 257-264 (with P. Nyikos).
28. "A note on removable pairs," **Proc. of the Fifth International Conference on the Theory and Applications of Graphs** (1991), 739-742 (with W.T. Trotter).
29. "A polynomial time approximation algorithm for Dynamic Storage Allocation", **Discrete Math.** **88** (1991) 231-237.
30. "Computing the dimension of N-free ordered sets is NP-complete", **Order** **6** (1989), 133-135 (with S.G. Penrice).
31. "The number of depth first searches of a poset", **Order** **6** (1989), 295-303 (with W. T. Trotter).
32. "Fibres and ordered set coloring", **J. of Combinatorial Theory Series A** **58** (1991), 158-164 (with D. Duffus and W. T. Trotter).
33. "Recent results on a conjecture of Gyárfás", **Congressus Num.** **79** (1990), 182-186 (with S.G. Penrice).
34. "The dimension of random ordered sets", **Random Structures and Algorithms** **2** (1991), 253-275 (with P. Erdős and W.T. Trotter).
35. "Radius two trees specify χ -bounded classes", **J. of Graph Theory** **18** (1994) 119-129 (with S.G. Penrice).
36. "Colorful induced subgraphs", **Discrete Math.** **101** (1992) 165-169 (with W.T. Trotter).
37. "On the game chromatic number of some classes of graphs", **Ars Combinatoria** **35** (1993) 143-150 (with U. Faigle, W. Kern, and W.T. Trotter).
38. "Long stars specify weakly χ -bounded classes", **Colloquia Mathematica Societatis János Bolyai** **60 Sets, Graphs, and Numbers** (1991), 421-428.

39. "On-line graph coloring", **On-Line Algorithms**, C. McGeoch and D. Sleator eds., (1992) 85-92 (with W.T. Trotter).
40. "An explicit 1-factorization in the middle of the Boolean Lattice", **J. of Combinatorial Theory Series A** **65** (1994), 334-342 (with D. Duffus and H. Snevily).
41. "First-Fit and on-line coloring of graphs which do not induce P_5 ", **SIAM J. on Discrete Math.** **8**, (1995) 485-498 (with S.G. Penrice and W.T. Trotter).
42. "On-line graph coloring and recursive graph theory", **SIAM J. on Discrete Math.** **7**, (1994) 72-89 (with S.G. Penrice and W.T. Trotter).
43. "Coloring interval graphs with first-fit", **Discrete Math.** **144** (1995) 47-57 (with Jun Qin).
44. "Recursive and on-line graph coloring", **Recursive Mathematics**, Elsevier, Y. Ershov, S. Goncharov, A. Nerode, and J. Remmel eds., to appear.
45. "Coloring planar graphs with an uncooperative partner", **J. of Graph Theory** **18** (1994) 569-584 (with W.T. Trotter).
46. "Applications of hypergraph coloring to coloring graphs which do not induce certain trees", **Discrete Mathematics** **150**, (1996) 187-193 (with V. Rödl).
47. "Some results on dimensions of certain families of ordered sets", **Order** **9** (1992), 103-110 (with Jun Qin and W.T. Trotter).
48. "The square of paths and cycles", **J. of Combinatorial Theory Series B** **63** (1995) 55-64 (with Genghua Fan).
49. "The dimension of suborders of the Boolean Lattice", **Order** **11** (1994) 127-134 (with Brightwell, Kostochka, and Trotter).
50. "Classes of graphs that are not vertex ramsey", **SIAM J. on Discrete Math.**, to appear.
51. "Hamiltonian square-paths", **J. of Combinatorial Theory B**, **67** (1996) 167-182 (with Genghua Fan).
52. "The order dimension of 1-sets versus k-sets", **J. of Combinatorial Theory A** **73** (1996) 219-228.
53. "On-line Coloring of Perfect Graphs", **Combinatorica** **16** (1996) 479-491 (with K. Kolossa).
54. "Game chromatic number and treewidth", submitted (with Z. Tuza).
55. "Square hamiltonian cycles in graphs with maximal 4-cliques", **Discrete Math**, to appear (with J. Quintana).

56. "Partitioning a graph into two square-cycles", **J. of Graph Theory**, **23** (1996) 241-256 (with Genghua Fan).
57. "Classes of graphs that exclude a tree and a clique and are not vertex Ramsey", **Combinatorica** **16** (1996) 493-504 (with Yingxian Zhu).
58. "The dimension of finite subsets of κ ", **Order**, **13** (1996) 227-231 (with E. Milner).
59. "Radius three trees in graphs with large chromatic number and small clique size", submitted (with Yingxian Zhu).
60. "On-line coloring k-colorable graphs", **Israel J. of Math**, to appear.
61. "Coloring graphs on-line", submitted.
62. "Interval orders and dimension", **Discrete Math**, to appear (with W.T. Trotter).
63. "Hamiltonian Chains in Hypergraphs", submitted (with Gyula Y. Katona).

Other References:

64. On-line Algorithms and Polynomial Time Approximation, Research proposal for Grant N0001490-J-1206 (1989).
65. On-line Algorithms and Other Combinatorial Optimization Problems, Research proposal for Grant N0001490-J-1206 (1992).
66. P. Erdős, Problem 9, **Theory of Graphs and Applications** (M. Fieldler ed.), Czech. Acad. Sci. Publ., Prague (1964) 159.
67. Z. Füredi and J. Kahn, "On the dimension of partially ordered sets", **Order** **3** (1986) 15-20.
68. A. Gyárfás, "On Ramsey Covering-Numbers", **Coll. Math. Soc. János Bolyai** **10, Infinite and Finite Sets**, North-Holland/American Elsevier, New York (1975), 801-816.
69. A. Gyárfás, "Problems from the world surrounding perfect graphs", **Zastowania Matematyki Applicationes Mathematicae** **XIX**, (1985), 413-441.
70. A. Gyárfás, E. Szemerédi, and Tuza, "Induced subtrees in graphs of large chromatic number", **Discrete Math.** **30**, (1980), 235-244.
71. N. Sauer, "Vertex partition problems", **Combinatorics, Paul Erdős Is Eighty (Volume I)**, (1993) 361-377.